

facts: About



The Use of Alcohol-based Hand Sanitizers in Ohio Schools

Summary

Alcohol-based hand sanitizers can be effective when used for hand hygiene. They may be safely and effectively used in the school setting if guidelines are followed.

Background

The Centers for Disease Control and Prevention (CDC) recommend hand washing to reduce infection transmission. Effective hand washing can be done with either soap and water or an alcohol-based hand sanitizer. Because of the widely recognized benefits of hand hygiene, Jarrod's Law (Ohio Administrative Code [OAC] 3701-54-09) requires all schools to develop and implement a written protocol for hand washing in each school. Such a protocol may require the use of soap and water and/or alcohol-based hand sanitizers.

Purpose

In general, if hands are visibly soiled, soap and water should be used for hand hygiene. In other circumstances, the use of alcohol-based hand sanitizers can be as effective as soap and water. In the case of younger students who may not thoroughly wash, rinse and dry hands, the use of alcohol-based hand sanitizers may be more effective than and preferable to the use of soap and water when hands are not visibly soiled. In addition, the use of alcohol-based hand sanitizers can allow a large number of students and staff to cleanse their hands in a time-efficient manner. Alcohol-based hand sanitizers must be at least 60 percent alcohol to be effective; the use of other types of cleansers is not recommended. Alcohol-based hand sanitizers are not effective in removing nut allergens from surfaces. It is recommended that unscented products be used in the school setting, as scented products may trigger asthma or allergy symptoms.

Guidelines for Use

Hand hygiene with an alcohol-based hand sanitizer should be done frequently, and especially at the following times:

Before

- Preparing food or eating
- Inserting or removing contact lenses
- Rendering first aid or treating a person who is ill
- Administering medication or completing a medical procedure

After

- Any contact with blood, body fluids or soiled objects
- Using the toilet
- Assisting with personal hygiene, such as changing diapers
- Touching animals or animal waste
- Food preparation, especially after handling raw meat, poultry or fish
- Blowing your nose, using a tissue or coughing or sneezing into your hands
- Treating or touching open wounds or a person who is sick or injured
- After handling trash or garbage
- After playing outdoors

In Ohio schools, all students and staff should receive instruction on the appropriate and effective use of hand sanitizers. The following recommendations are for best practice in the use of alcohol-based hand sanitizers in Ohio schools.

1. Alcohol-based hand sanitizer dispensers should be located in places where students are supervised by school staff at all times
2. Do not allow students or staff to ingest alcohol-based hand sanitizers
3. Alcohol-based hand sanitizers are flammable-do not use while smoking or near open flame

The use of Alcohol-based Hand Sanitizers in Ohio Schools

4. It is recommended that the school purchase alcohol-based hand sanitizers for use in each classroom and cafeteria, just as it purchases soap and towels for restrooms and kitchens. The advantages for such a school purchase include:
 - a. The school will have a material safety data sheet (MSDS) for all alcohol-based hand sanitizers in the building
 - b. Refills for the alcohol-based hand sanitizers can be stored with cleaning supplies away from heat sources or open flames and in accordance with Ohio Fire Code regulations (OAC 1301:7-7-3405.5)
5. Alcohol-based hand sanitizer dispensers must be located so they do not drip liquid on the floor to create a fall hazard. A “drip catcher” may be placed on the dispenser to prevent this

Instructions for Placement and Storage

The placement of the alcohol-based hand sanitizers, if wall mounted, should be consistent from room to room and in compliance with the Ohio Fire Code regulations (OAC 1301:7-7-3405.5) summarized here. Dispensers must:

- Have a maximum capacity of 68 ounces
- Be a minimum of 48 inches apart
- Be installed in locations that are not above, below or directly adjacent to electrical outlets, switches, appliances, devices or any other potential ignition source
- Be placed between 42 and 48 inches from the floor
- Be manually activated
- Be properly supplied with sprinkler systems if placed in carpeted areas

In addition, alcohol-based hand sanitizers in school hallways or corridors must comply with Ohio Fire Code regulations (OAC 1301:7-7-3405.5) summarized here:

- There can be no aerosol containers in the hallway
- Dispensers can have a maximum capacity of 41 ounces
- There can be no more than 10 gallons in the hallway
- The hallway must be a minimum of 72 inches in width
- There can be no unapproved projections into the hallway

References

APIC. (2005.) *Prevention Experts Stress Importance of Hand Hygiene Compliance*. Available online: <http://www.apic.org/AM/Template.cfm?Section=Search§ion=Regulatory1&template=/CM/ContentDisplay.cfm&ContentFileID=1458>

CDC. (2008.) *Clean Hands Save Lives!* Available online: <http://www.cdc.gov/cleanhands/>

CDC. (2002.) *Hand Hygiene Guidelines Fact Sheet*. Available online: <http://www.cdc.gov/od/oc/media/pressrel/fs021025.htm>

CDC. (2008.) *Hand Hygiene Saves Lives*. Available online: <http://www.cdc.gov/handhygiene/>

CDC. (2006.) *Hand Sanitizer Alert*. Available online: <http://www.cdc.gov/ncidod/eid/vol12no03/05-0955.htm>

CDC. (2007.) *MMWR Hand-Washing Recommendations to Reduce Disease Transmission From Animals in Public Settings*. Available online: <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5605a4.htm>

Jarod's Law.
Ohio Administrative Code 3701-54-09 (M) (10).
Available online: <http://codes.ohio.gov>

Mayo Clinic. (2008). *Hand washing: An easy way to prevent infection*. Available online: <http://www.mayoclinic.com/health/hand-washing/HQ00407>

Ohio Fire Code.
Ohio Administrative Code 1301:7-7-3405.5.
Available on page 394 online: https://www2.iccsafe.org/states/ohiofire/OH_Fire/fire_frameset.html